

Athlete Report





Welcome to your unique, personalised athlete DNA report!

DNA sequencing has opened the door to personalised approaches to health and fitness, enabling a more intelligent approach to training, recovery and performance. In this report we present elements of your unique DNA profile that have been shown to affect athletic performance. It is well understood that athletic performance depends on interactions between genetic and environmental factors. Your genes are not your destiny. By aligning your training, nutrition and sleep practices with your genetics you can realise your athletic potential.

Whether you are an elite athlete, a 'weekend warrior', or just want to get healthier, your athlete DNA report will enable you to:

01 TRAIN

tailor a programme that will be most effective for you to support cardio-vascular adaption, strategic fuelling and structural strength

03 PERFORM

balance stress and stimulation when it matters most - to achieve your personal best

02 RECOVER

build resilience and reduce your risk of injury and illness by controlling inflammation, reducing oxidative stress and optimising sleep

LGX

The nutritional genomics team at Lifecode Gx are experts in interpreting DNA results, examining the most up to date research in the field and distilling it into relevant, meaningful and practical advice.

To get the most benefit from this report we recommend that you work with a qualified sports nutrition professional.



Nadura's functional sports performance programme focuses on the biochemical individuality of their athletes. Providing a guided diet and supplements plan, personalised to their genetic make-up, biochemistry and physiological training requirement.

This allows their expert nutritionists to maximise results for both professional and amateur athletes alike.

How to Read Your Report - Your Results in Detail

Genes

Genes are listed in alphabetical order, by name. The short name (e.g. VDR) is usually the acronym of the long name (e.g. Vitamin D Receptor). The description explains the gene function (purpose).

VDR Vitamin D (1,25- dihydroxyvitamin D3) Receptor

Vitamin D receptors (VDR) are located in almost every tissue within the human body. Acting via VDRs, vitamin D impacts a wide variety of health and performancerelated variables, such as exercise-induced inflammation, tumour suppressor genes, neurological function, cardiovascular health, glucose metabolism, bone health and skeletal muscle performance.

rs1544410 TC V Potential lower response to Vitamin D and increased need To reduce risk of stress fractures and shin splints, and muscle fatigue and scieness ensure sufficient Vitamin D intake and UV exporure. Training should include weight bearing exercises. Dietary sources of vitamin D include oily fish - salmon, mackerel, sardines and tuna, milk and egg yolks.

Personalised Result

Highlight Colour

The genotype result highlight indicates the potential effect of the SNP on gene function in a particular context.

RED the effect of the variant is negative

- **AMBER** the effect of the variant is neutral (no effect) or is subjective
- **GREEN** the effect of the variant is positive

Arrow Direction

The direction of the arrow indicates the potential effect of the SNP on gene expression - it can increase or decrease activity, or neither.

The 'rs' number is the reference sequence number that

Your genotype result is shown as two letters (A,G,T or C) which represent the DNA bases present at that location.

identifies a specific location on the genome. It is also

known as a SNP (Single Nucleotide Polymorphism)

pronounced 'snip', polymorphism or mutation.

- ▲ up-regulates or increases the activity and effect of the gene
- down-regulates or decreases the activity and effect of the gene

No arrow - no effect on the activity of the gene



By understanding your genetic makeup you can tailor a training programme that will be most effective for you.

Cardiac Output

ACE

AGT

Your cardiac output potential is **elevated**. This can be beneficial for sports that require high-intensity power such as sprinting or weight-lifting. However, high sodium levels and fluid retention can elevate blood pressure.

Monitor and control for high blood pressure. Potassium, magnesium and calcium can moderate the blood pressure raising effects of sodium (salt). Maintain a healthy body weight and exercise regularly.

Blood Circulation

BDKRB2 NOS3 VEGFA

Your circulation potential is **normal**.

Aerobic exercise will stimulate blood vessel growth and improve circulation of oxygen and energy needed for training. Increasing intake of foods rich in nitric oxide, such as beetroot, can improve blood vessel growth.

Cardio

The cardiovascular system is essentially made up of two parts - the heart ('cardio') and the blood vessels ('vascular' or 'circulation'). Its primary functions are to transport nutrients, hormones, gases, and waste to and from our cells and to regulate body temperature and fluid balance.

Exercise places a greater demand on cardiovascular functions. As working muscles require more oxygen and nutrients than normal, they produce more waste products and generate more heat. The good news is the body is designed to adapt to these demands. The extent of adaptive response is determined by training, lifestyle, and genetic predisposition.

Although the cardiovascular system is integrated, we present the genes and variants according to their impact on the 'cardio' or 'vascular' components - described in terms of cardiac output and circulation.

Cardiac Output

Cardiac Output is the amount of blood pumped by the heart, measured in litres per minute. It ranges from 5 l/min at rest up to 40 l/min during maximal effort. Cardiac output increases during exercise in order to meet demands for more oxygen and nutrients and removal of waste. It has two components:

• Heart Rate is the number of times the heart beats per minute (bpm). A normal resting heart rate range is between 60-100bpm.

• Stroke Volume is the volume of blood pumped by the left ventricle with each heart beat.

Blood Pressure

Blood Pressure is a measure of the force being exerted on the walls of arteries as blood is pumped out of the heart. Systolic (the top number) is measured as the heart contracts, and diastolic (the bottom number) as it relaxes between beats. A blood pressure reading of 120 (systolic)/ 80 (diastolic) measured in mmHg is considered normal.

High blood pressure (hypertension) occurs when resting systolic blood pressure readings consistently exceed 140mmHg and/ or diastolic readings exceed 90mmHg. Increases in cardiac output, blood volume, blood viscosity (thickness) or systemic vascular resistance can all increase blood pressure.

Low blood pressure (hypotension) and low heart rate are very common amongst athletes, and are usually a sign of good cardiovascular efficiency.

Cardio Genes

The **ACE** and **AGT** genes form part of the reninangiotensin-aldosterone system (RAAS) which plays an important role in regulating blood volume and blood pressure.

Variants on the ACE and AGT genes can increase RAAS activity, including levels of angiotensin II. This has multiple potent effects which can contribute to raised blood pressure:

- Constricting (narrowing) blood vessels increasing blood pressure directly.
- Triggering the release of aldosterone, causing the kidneys to retain sodium and water and excrete potassium increasing blood volume.
- Increasing thirst and appetite for salt which can also lead to increased blood volume.

Higher RAAS activity is also associated with greater muscle growth (hypertrophy) and left ventricular size which has been linked with elite status in anaerobic/ high-intensity sports.

Potassium is an essential nutrient that can counter-balance the blood pressure raising effects of sodium (in salt).

Apricots, bananas and coconut water are good natural sources of potassium and easy to eat on the go!

Circulation

The circulatory, or 'vascular' system, is made up of vessels that carry blood and lymph through the body.

Arteries carry oxygen rich blood away from the heart. The arteries break down into smaller and smaller branches, called capillaries. As blood moves through the capillaries, the oxygen and other nutrients move out into the cells, and waste matter from the cells moves into the capillaries. Blood is then moved from the capillaries to the veins, which carry it back to the heart.

The circulatory system directs blood to where it is needed most. During exercise blood flow is directed away from core systems and organs to skeletal muscle. The body adapts to exercise by growing new blood vessels - this is called 'angiogenesis'.

Circulation Genes

As its name suggests, vascular endothelial growth factor (coded for by the **VEGFA** gene) promotes the growth of new blood vessels. VEGF also stimulates endothelial nitric oxide synthase (**eNOS**) which supports the production of nitric oxide from l-arginine. Nitric oxide relaxes and dilates blood vessels, thereby increasing blood circulation.

Variants on the **BDKRB2** gene increase sensitivity to bradykinin, lowering RAAS activity and blood pressure and increasing circulation. Athletes with lower RAAS activity and good circulation tend to perform better at endurance sports.

> The body can manufacture **nitric oxide** from the amino acids I-citrulline and I-arginine with the help of the eNOS gene.

Dietary nitrates can also be turned into nitric oxide via a separate oxygen-independent pathway. Beetroot juice has a particularly high nitrate content.

ACE Angiotensin I Converting Enzyme

Angiotensin I Converting Enzyme is able to cleave (cut) proteins. By cutting angiotensin I at a particular location it converts it to angiotensin II.

Angiotensin II causes blood vessels to constrict (narrow), raising blood pressure. As a growth factor, it is associated with greater muscle volume and strength which may be beneficial to high intensity (power) oriented sports.

rs4343 AA

Normal (not increased) ACE activity. Not predisposed to water and salt retention or high blood pressure.

More likely to excel at, and benefit from, low-medium intensity/ cardio oriented activities rather than power oriented. Ensure sufficient hydration and electrolyte intake.

AGT Angiotensinogen

Angiotensinogen is the primary controller of the reninangiotensin-aldosterone system (RAAS) which regulates blood pressure and the balance of fluids and salts in the body.

Variants are associated with higher activity and more rapid conversion to angiotensin I and II. Although higher angiotensin II is associated with higher blood pressure and sodium retention, it is also linked to increased cardio output which is beneficial for power (high-intensity) performance.



Increased AGT activity and angiotensin II levels. Mixed impact - greater cardiac strength (positive) however water and salt retention may elevate blood pressure (negative).

Likely to excel at, and benefit from, power (high intensity) oriented activities. Monitor blood pressure and ensure adequate potassium intake to balance the effects of sodium.

BDKRB2 Bradykinin Receptor Beta 2

Bradykinin causes blood vessels to dilate (widen), and blood pressure to fall. It also stimulates uptake (storage) of glycogen by skeletal muscle.

Variants on the Bradykinin Receptor Beta 2 gene increase sensitivity to bradykinin, so less bradykinin is needed to stimulate vasodilation, and improve blood circulation.



No variance. Normal (low) bradykinin sensitivity. You may need higher levels of bradykinin for its effects to be felt.

Aerobic exercise can raise bradykinin levels and help to lower blood pressure and improve circulation.

Some pages have been omitted in the sample

Phosphocreatine

MTHFR

Your potential to generate a lot of energy, quickly, is **good**. This is beneficial for high intensity activities that require short bursts of energy such as sprinting or weightlifting.

MTRR

To maximise your potential, ensure adequate intake of creatine (red meat and fish are good natural sources). Vitamins B9 and B12, magnesium and zinc will help to support methylation.

Glycolysis

ADRB2

Your potential to generate energy anaerobically from glucose is **elevated**. This may be advantageous for speedoriented activities such as short distance swimming or sprinting.

Excess glucose will be stored as glycogen or fat. To avoid weight gain, limit your intake of simple carbohydrates.

Aerobic

AMPD1 PPARA ▼▼

l GX

PGC1A

Your potential to adapt to using fat for fuelling endurance activities is **reduced**. This can be a disadvantage for sports that require sustained energy release such as long distance running or cycling.

Good sources of healthy fat include avocados, nuts, seeds, fatty fish and organically farmed meats.

Fuel

The body uses a mixture of carbohydrates, fats and proteins as sources of energy. The balance at any point in time depends on intensity of activity, availability of fuel type, and genetics.

Carbohydrates are the favoured fuel source as they are quickly and easily converted, providing immediate energy for high-intensity activities. Fat takes longer to convert and supplies energy to support low to moderate intensity exercise. Protein can provide energy when other sources are scarce.

There are three systems for generating energy, each of which has a different biochemistry and rate of ATP production:

- Phosphocreatine
- Glycolysis
- Aerobic

Phosphocreatine

The phosphocreatine system uses creatine to supply a phosphate group to recycle ADP to ATP and release energy. It is able to generate energy quickly, but only for a short period of time. This is useful for high-intensity activities such as weight training or sprinting.

Creatine synthesis can be impacted by variants on the **MTHFR** and **MTRR** genes and supply of vitamins B9 (folate) and B12. Creatine can also be obtained directly from the diet.

Creatine synthesis is heavily dependent on methylation and uses up significant amounts of a substance called SAMe. Variants on the MTHFR and MTRR genes can also impact SAMe levels.

To support methylation you need a good supply of B vitamins including B9 (found in green leafy veg, citrus fruits, nuts and beans) and B12 (found in meat, eggs and fish).

Glycolysis

Glycolysis is the break down of glucose to generate energy. It can be anaerobic or aerobic. Anaerobic glycolysis is the major source of energy during strenuous exercise without the use of oxygen.

Glycogen is broken down into glucose and rapidly converted to ATP. However, as each glucose molecule only produces a small amount of ATP, it is inefficient. After a few minutes of exercise, the body starts to switch over to the aerobic system.

Variants on the **ADRB2** gene can affect the amount of blood glucose available to fuel glycolysis.

Aerobic

When oxygen is available, the body is able to generate ATP by breaking down carbohydrates and fat aerobically.

Initially, most energy production is fuelled by muscle glycogen. After about two hours of high-intensity exercise, the fuel source switches to carbohydrate and fat (lipolysis). Although fat is energy dense its conversion to ATP is slow compared to carbohydrate.

Variances on **AMPD1**, **PGC1A** and **PPARA** genes affect your aerobic capacity and adaptability to utilise fat as fuel alongside carbohydrate.

Fat adaption is the process of training your body to break down fat for energy. As well as improving endurance ability, this can have many health benefits including weight loss, balanced energy levels and cardiovascular health.

Oily fish, avocados, olive oil and coconut oil are good sources of healthy fats.

ACTIG Algha 3 Actinin

The ACTED game emcodes, applies action 3 which a present in type 1 (feet twistif) forms in destend muscle and is responsible. For generating force at high whichly t is collectually involve as the 'generating process'.

Fing House Annual Chard Languages (et al., 2011) is relationed to an IREPTR Descention the contained T administration at the container (etc). By Section 2 the Same Seatistic Millions, there is a many potential for desceletaring situation basis. In Biome, Manual, 1995, New Sectional Administration of the genes is associated with other emburgence attribute status.

101815738

and the second



CDL1A1 Collagen, Type 1, Alpha-1

the Ensure addresses include of alignment's co-



and the set of the set

117546410

natio, variancia, fearmar gassel, sensitivity to = 0.

To recluce risk of green Tractures and stor splitting, and muscle fargue and screeness amount sufficient interest 0 make and UV exposure. Detary sources of etamore 2 millude obj fish - salmon, machinet, sandtrees and sing, with and app sole. Some pages have been omitted in the sample

Muscle

ACTN3

Your muscle fibre type is **fast twitch** dominant. This is the 'speed' version of the gene which is beneficial for sports that require quick bursts of power, such as sprinting.

You can increase the size and strength of fast twitch muscle fibres, and power capacity, by regular strength training, however you ability to develop slow twitch muscle fibres may be limited by your genetic make-up.

Connective Tissue

COL1A1

Normal (healthy) collagen synthesis. Your risk of tendinopathy type injuries is **not elevated**.

Your diet should include collagen rich nutrients, including vitamins C, E, manganese and zinc.

Bone

VDR

l GX

You are able to utilise vitamin D and vitamin K **effectively**. The body needs vitamin D, vitamin K, calcium and phosphorus to build strong bones.

VKORC1

Your body can make vitamin D directly if you have sufficient exposure to sunlight. Vitamin K is abundant in leafy green vegetables, brussels sprouts, cabbage and broccoli. Weight bearing exercise will also help to improve bone strength.

Structure

The musculoskeletal system gives the body structure and stability and enables movement. It is made up of bones, skeletal muscle and connective tissues - tendons, ligaments and cartilage.

Whilst genes may determine our physical limits, we all have some potential to alter aspects of our size and strength through physical training and lifestyle choices. Although they are interconnected, we look at genetics affecting the following three components:

- Muscle
- Connective tissue
- Bone

Muscle

Skeletal muscles are made up of a combination of two main types of fibre, called slow-twitch (type I) and fasttwitch (type II), which influence how muscles respond to training and physical activity.

Slow-twitch muscle fibres can work for a long time without tiring so are good for endurance activities, like long distance running. Fast-twitch muscle fibres fatigue more quickly but support rapid, bursts of movement needed for power activities, such as sprinting or weightlifting.

Humans are genetically predisposed with a larger percentage of one type of muscle group over another. The **ACTN3** gene has a significant influence on the number of type II (fast twitch fibres).

Although the number of muscle fibres cannot be increased, cells can be induced to grow larger through hormone signalling and strength training, supported by high quality protein intake.

Amino acids, the building blocks of proteins, are needed to build and repair muscle tissue, to increase strength and reduce soreness. Nine of these amino acids are categorised as 'essential', as the body cannot make them. The complete range of amino acids can be sourced in one hit from animal proteins (meat, fish, eggs) or by consuming a variety of plant-based foods.

Connective Tissue

Collagen is the main structural protein in connective tissue and the most plentiful protein in the body. It is mostly found in fibrous tissues, tendons, ligaments and skin, and in bone. It also constitutes a small proportion of muscle tissue.

There are more than 30 genes associated with synthesis of collagen, each coding for a different type. The **COL1A1** gene codes for type I collagen, the most abundant form, found in tendons, skin, artery walls, muscles, bones and teeth.

Collagen is made from amino acids (principally, glycine and proline) and requires vitamin C as a co-factor. Longterm deficiency in vitamin C results in impaired collagen synthesis and scurvy-like symptoms. **Vitamin C** is essential for the formation of collagen. Athletes may be at greater risk of vitamin C deficiency due to the wear and tear associated with training.

Good sources include citrus fruits, kiwi, strawberries, broccoli and tomatoes.

Bone

Bones are made from a matrix of collagen into which bone building minerals - calcium, phosphorus, magnesium and potassium are deposited. Bones are in constant flux, endlessly being broken down (by osteoclasts) and re-created (by osteoblasts).

Weight-bearing exercise combined with sufficient protein, mineral and vitamin intake stimulates bone growth. Vitamins D and K work together to support the absorption and distribution of calcium - increasing bone formation and preventing bone loss. Sensitivity to vitamin D can be impacted by variants on the Vitamin D Receptor (**VDR**) gene. Variants on the **VKORC1** gene can impact your ability to convert vitamin K to its usable form.

Vitamin D

Vitamin D is called the 'Sunshine Vitamin' because the body can make its own vitamin D when skin is exposed to sunlight.

Once in the body, vitamin D is converted to the activated form of vitamin D3 (also called calcitriol, or 25(OH)D)) - which acts via the vitamin D receptor (VDR).

Vitamin D is essential for the maintenance of bone mineralisation through the regulation of calcium and phosphorus levels. Deficiency can increase the risk of lower bone density - broken bones, shin splints and osteoporosis. Other symptoms of vitamin D deficiency include muscle weakness, difficulty thinking clearly and unexplained fatigue.

Sun exposure, diet, skin colour, age and body weight, have varied and substantial impacts on vitamin D levels. Although sunlight is the best source, small amounts of vitamin D are available in food - mackerel, salmon, tuna and sardines are particularly good sources.

Vitamin D supplements are widely available, however you can have too much of a good thing. Vitamin D toxicity can cause abnormally high blood concentrations of calcium, leading to over-calcification of bones, soft tissues, heart and kidneys. Bone is a living organ that is continuously being reshaped. **Vitamin D and vitamin K work together** to support absorption and distribution of the calcium needed to build strong bones.

Vitamin K can be found in dark green, leafy vegetables and brassicas (broccoli, brussels sprouts and cabbage). Oily fish, such as mackerel, sardines, salmon and tuna are good sources of vitamin D.

A vitamin D blood test is the only way to categorically determine if you are deficient in vitamin D.

ACTING Algibra 3 Actiniin

101012738

In the second second





CDL1A1 Collagen, Type 1, Alpha 1

A contact on the light region of the CD. We prove the the second of second second second second and the second sec

of the series reported that of converting

and the second se from series a because and the second second second our streng hards an even on even and



Some pages have been omitted in the sample



Recovery is a major factor in developing strength and endurance. It is as important as the training itself.

 \mathbf{P}

Inflammation

IL6 ▼

Your predisposition to exercise induced inflammation is **reduced**. You may be less susceptible to muscle damage and soreness in response to training and benefit from shorter recovery times.

TNF

Ensure adequate recovery times and follow an anti-inflammatory diet including omega 3 fatty acids.

Inflammation

Athletes who don't recover adequately from workouts become vulnerable to injury or illness. To help prevent these problems the body relies on its natural inflammatory and anti-inflammatory mechanisms.

When injured, the body creates natural inflammation around a wound, making it reddish, swollen and hot. This is called acute inflammation, and is the body's way of recovering from the daily wear and tear of physical and chemical stress.

Chronic inflammation, on the other hand, develops when the body is unable to control the inflammatory mechanism. This is often due to nutritional imbalances, stress, overtraining or insufficient sleep.

Although a certain amount of muscle damage is needed to promote adaptions to muscle size, strength and efficiency, excessive damage or inadequate recovery can increase inflammation, muscle fatigue and risk of injury. As we get older we are likely to experience more muscle damage and take longer to recover.

Exercise reduces expression of inflammation-promoting cytokines, such as tumour necrosis factor (**TNF**) and promotes the expression of the anti-inflammatory cytokine interleukin 6 (**IL6**).

Gene variants that influence inflammatory response have been linked to differences in exercise induced muscle damage and recovery times.

C-reactive protein (CRP), which is secreted by the liver, circulates in the bloodstream and is a commonly used **biomarker for inflammation**.

Pro-inflammatory

TNF is a pro-inflammatory cytokine that plays a vital role in the body's immune response. However, overproduction of TNF can result in chronic inflammation and autoimmune and immune-mediated disorders.

The inflammatory effects of TNF are inhibited by several natural compounds including curcumin (found in tumeric), catechins (in green tea), omega 3 fatty acids (in oily fish) and echinacea. More recently, exercise has also been shown to blunt the effects of TNF.

Anti-inflammatory

Plasma IL6 levels increase exponentially (up to 100 fold) with exercise depending on intensity and duration, muscle mass, and endurance capacity.

IL6 increases ATP generation, induces lipolysis and fat oxidation and maintains blood sugar homeostasis. It also has strong anti-inflammatory effects including inhibition of TNF activity and has consistently been shown to protect against muscle damage.

Knowing how someone is likely to respond to a particular type of exercise could help practitioners individualise the exercise training of their athletes; maximising recovery and reducing the risk of overtraining. The Omega-3 fats found in fresh, 'oily' fish such as salmon, sardines and anchovies, have powerful **anti-inflammatory** effects.

Ginger, turmeric, citrus peel and foods in the onion family (shallots, chives and garlic) all have anti-inflammatory effects.



ACTINI Alighta 3 Activity

The ACTED game emcludes applies actions 3 adults a present in type 1 (fast tasks)) films in destinat multicle and is responsible. For generating force at high selectly tis collespoid, income as the 'generat' gene.

The most instantic had variant on A, Thill L referenced to an ADTTR because the variant 1 alters excludes a true color (2). By lending the fact testich flores there 5 most potential for developing time testich flores. Hence, this room functional attracts of the gene 5 accordent with effect modulations attracts for the

101815738

and the second second





CDI,1A1 Collagen, Type 1, Alpha-1

111546410

. .

to genetic variance. Normal good, sensitivity to Hamme 3

To reduce risk of green fractures and drive splitting, and multicle fargues and screeness emure sufficient vituence. Dimple and UV exposure. Descry sources of elument Dimple only fish - salmost, machinet, sandther, and ture, risk and egg unles.

Oxidative Stress

GSS NOS3 SOD2 ▼

Your genetic predisposition to oxidative stress is **normal**. This means your susceptibility to cellular damage caused by exercise is not elevated.

Antioxidants protect cells against free radicals and help repair cellular damage. Ensure your diet is rich in antioxidants including vitamins A, C and E, found in many fruits, vegetables, nuts and some fish, and meats.

Oxidative Stress

Free radicals are a normal by-product of the body's energy-generating biochemical processes. They are highly reactive with other molecules, and can damage DNA, proteins and cellular membranes. The balance between oxidation and anti-oxidation is critical to maintaining healthy biological systems.

Dietary anti-oxidants such as vitamin C, vitamin E, carotenoids and polyphenols are free radical scavengers that interact with free radicals to deactive them. However, the major role in anti-oxidant defence is fulfilled by the body's own 'master antioxidant', glutathione. Oxidative stress can be managed by addressing both cause and effect.

Causes of oxidative stress can be psychological, emotional and social, as well as physical. Activities like meditation, walking in nature and yoga can help to relieve the sources of some of these stressors.

As an athlete, a certain amount of oxidative stress due to physical activity is inevitable. Individuals who have variants on the Nitric Oxide Synthase (**NOS**) or Superoxide Dismutase (**SOD**) genes may be more susceptible to oxidative stress than others.

Increasing intake of antioxidant foods and herbs will help improve antioxidant status and reduce the effects of oxidative stress. However, the most prolific antioxidant in the body is glutathione, which is made from the amino acid cysteine, with help from the Glutathioine Synthetase (**GSS**) gene.

Glutathione is the most prolific antioxidant in the body. It is often referred to as the 'master antioxidant'.

The amino acid, **cysteine** is needed to make glutathione. Cysteine is found in high protein foods, including meat, eggs and dairy, and in lesser amounts in sulphur containing foods, such as broccoli, Brussels sprouts, peppers, onions and garlic.

Foods High In Antioxidants:

Goji berries, Blueberries, Cranberries, Blackberries, Dark chocolate, Pecan nuts, Coriander, Kidney beans

Herbs High In Antioxidants:

Basil, Cayenne pepper, Cloves, Cinnamon, Cocoa, Garlic, Ginger, Green tea, Oregano, Parsley, Turmeric, Thyme

ACTING Alights 3 Actinity

101012738

and the second second

AND INCOME.



CDI, 1&1 Callagen, Type 1, Alpha-1

the state has been a set of the s

A contact on the light region of the CCL will prove have tioned in the set of print the set of the set rear and had been read of an experiment.

and the second second second

with the two of the second second second second

Passing and a the second of the second second the same in the last of the same the second se

117548470

Sleep

CLOCK **V**

The **in-betweener** may find it easier to adjust their sleep and wake routine to fit their environment.

PER1

Try to maintain a regular sleep routine and avoid stimulants and bright light close to bed time.

Sleep

Sleep is vital to overall health and enabling repair and recovery from exercise. Like other aspects of biology sleep is individual - sleep and wake patters are influenced by genetics.

The term chronotype describes someones sleeping characteristic - if they are naturally a morning person (early bird) or evening person (night owl) or, the most frequent type, an 'in-betweener'. Understanding someones chronotype can enable sleep patterns to be adjusted to optimise recovery and performance.

Genetic testing can indicate which chronotype - 'earlybird', 'night-owl' or 'in-betweener' - an individual is most likely to associate with.

An early birds body clock runs slightly fast, and their circadian rhythm is just short of 24 hours. They find it easier to wake naturally and focus and perform better in the morning. Early birds are less likely to feel tired in the day and go to bed reasonably early.

A night owls body clock runs slightly slow, and their circadian rhythm might be twenty four and a half hours long. They find it difficult to wake and get up early and are more likely to peak around late afternoon or early evening.

The majority of people are 'in-betweeners' with a circadian rhythm just over or just under twenty four hours. Almost no-one has a 'perfect' twenty four hour cycle!

If you know your chronotype you may be able to schedule activities that require most focus and energy to coincide with your 'peak' performance time. Alternatively you can adjust your circadian rhythm by adopting regular sleep and wake times. Having a **regular wake time** is one of the most effective ways of re-training your body clock. If you are tired from getting up early, and exercising, your 'going-tosleep' time, should adjust naturally.

Foods that contain tryptophan can help the body make the sleep hormone,

melatonin. Good sources include bananas, almonds and turkey.

ACTIG Algha 3 Actinin

The ACTED game emcodes alpha action 3 adults a present in type 1 (fast tasks)) forms in desteral muscle and a responsible for generating force a high estimaty (a colloquially broast as the types) gene.

The most researched variant on ACMUS Loddened to a MOTH because the variant 1 alere encodes a singcolor (0). By lending the fast testsch filmes there 5 more potential for another statistic filmes there has potential for another of the gree 5 essected with eller endurance athlete status.

141815738





CDL1A1 Collagen, Type 1, Alpha-1

States and a



n genetic senancis. Hormat good senatory t Name 3

Schoolsander insk of appendix Nachduress and altern splitters, and multicle falligues and scoreneess ensure sufficient visuation (1) multicle and UV exposure. Destany solutions of eligenees 2) multicle obj. Roll. Saltman, machinest, sandtrees and surgationsk and egg unles.



Find the perfect balance to acheive your personal best.

[

Balance

5-HT2A LCT

Your susceptibility to symptoms of GI (gastrointestinal) distress is **normal**. However, you may experience symptoms in response to generic and specific triggers. For some people, foods that contain gluten, dairy or tryptophan (which can raise serotonin levels), may trigger adverse GI responses.

HLA-DQA1

Balance

The last thing you want to affect your performance on the big day is stomach problems, or worse! However, gastrointestinal (GI) symptoms, are common in athletes, and certain individuals or groups (elite athletes, women and endurance athletes) are more likely to be affected than others.

The impact of high-intensity physical activity on gut blood flow, motility and permeability can affect anyone. However, there are some dietary triggers that affect people quite differently and are reliably linked to genetics.

Athletes are more likely to experience symptoms relating to GI (gastrointestinal) damage and distress than nonathletes. Whilst mechanical effects, such as bouncing of organs, are difficult to avoid, habitual aspects, including nutrition and supplementation, can be pro-actively managed.

Genetic variances can significantly impact an individual's response to specific nutrients, or food groups, such as gluten and dairy.

Whilst most babies can digest lactose, the sugar found in milk and other dairy products, a variant near to the **LCT**, lactase, gene determines if you can retain this ability as an adult. If you have the 'wild'/default version of the gene you are unlikely to be able to digest lactose. In this case, the mutated version of the gene is beneficial.

The **HLA** gene variant that can lead to celiac disease is thankfully quite rare. However, many people develop non-celiac gluten sensitivity which means eating gluten, found in wheat and other grains, can cause GI damage.

Whilst serotonin is commonly known as the 'happiness' neurotransmitter, it can also increase nervousness and anxiety, and gut motility. Serotonin levels can be adjusted by eating tryptophan rich foods such as almonds, white meats and dairy products. Many brands of **dietary supplements** include buffering agents, or 'fillers', such as sodium bicarbonate which may cause Gl irritation with no nutritional benefit.

Symptoms of GI distress or injury include, abdominal pain and/or distention, nausea, flatulence, belching, constipation, loose stools, heartburn, acid reflux, hunger pains or stomach rumbling.

Non-steroidal anti-inflammatories

(NSAIDs), such as ibuprofen, are amongst the most commonly used drugs worldwide, and their pain-relieving and antiinflammatory effects are thoroughly accepted. However, they can also cause GI perforation and bleeding. Risk increases with concurrent use of other drugs, including aspirin, alcohol use and H Pylori infection.

ACTING Alights 3 Actinity

101012730

and the second second



CDL1A1 Collagen, Type 1, Alpha 1

provided, repairing inclusions for the providence of the A Transition

A contact on the light region of the CCA. man and had been made of a property

- page series

the same is the man per second real of period want has ever express that if several has - We have a for manyou the real of

where the longers advectance relates of strategies (to

- and the party of the second a second second second second the second s started and the second provide the started of

137548410

Caffeine Sensitivity

ADORA2A

Increased sensitivity to caffeine. Likely to experience greater adrenergic effects increases in heart rate and improved mental focus - with relatively low dosage. High caffeine intake may cause feelings of anxiety and jitteriness and negatively affect performance.

Caffeine Metabolism

CYP1A2

Faster metabolism of caffeine may shorten the duration of its effect.

Motivate

To optimise performance on competition day you need to be physically and mentally on top form at the moments when it matters most. Many athletes achieve this with the help of caffeine. Caffeine has proven benefits including alertness, reaction times, concentration and endurance capability. It is used extensively in sport as a legal and safe performance booster.

Caffeine acts mainly on the central nervous system, stimulating adrenaline release and the classic 'fight or flight' response; and increasing sensitivity to dopamine. It also has a metabolic effect, increasing energy expenditure. The overall effect of caffeine is reduction of fatigue, and improved ability to tolerate stress and pain.

Now, with the benefit of genetic testing, it is possible to further personalise the use of caffeine.

Genetic testing can enable personalisation of timing and dosage of caffeine to optimise performance. An individual's sensitivity to caffeine can be impacted by which variant of the **ADRB2** and **ADORA2A** genes they carry. Variants on the **CYP1A2** gene will impact the time it takes for caffeine's effects to wear off.

The ADORA2A gene controls the response to adenosine which acts to block the effect of dopamine. By inhibiting ADORA2A, caffeine enables dopamine to exert its effects, such as mood enhancement, better memory, and feelings of pleasure.

When caffeine triggers the release of adrenaline, the adrenaline receptor, ADRB2, stimulates increases in heart rate, blood pressure, muscle strength and blood sugar as a means to get you ready for strenuous activity, or 'flight or fight'.

CYP1A1 is the main deactivator, or detoxifier, of caffeine. Variants on this gene, are reported to increase its activity and metabolise caffeine more quickly.

To get the best results from caffeine, consider environmental aspects - body size, habitual caffeine intake, and timing and duration of the event, alongside genetic analysis. Adrenaline can increase focus so much that **pain is forgotten**.

Coffee, tea, citrus fruits, bananas, chocolate, cocoa and vanilla can all **raise adrenaline levels**.

Take care not to overdo it, particularly if you have the more sensitive version of the ADRB2 gene.

ACTIG Algha 3 Actinin

The ACTED game emcodes applies actions 3 which a present in type 1 (fast twist)/1) forms in desting? muscle and is responsible for generating force a high estimate 1.1 collectually income as the 'queed' gene.

The most installant had variant on AC field is relationed to an BUTTR because the sariant T also encoded in a trop color (0). By limiting the fair testsch filmers there is more protection for developing time testsch filmers filmers, this room functional atmost of the gene is accorded with other enducance athlets totals.

101815738

and the second second



CDL181 Collagen, Type 1, Alpha-1

of the local division in the

- acception

> non-transmission and and the angles, they non-transmission adequate make of algebra (to contain a second

117546410

His protect carbonal, Horman genetic sensitivity to Hamme 2

To reduce risk of greens fractures and shor splitting, and muscle fatigues and screeness around sufficient vituation 0 multiple and UV exposure. Destary sources of etamore 2 millude obj fath - salmon, machinet, sandmes and turus - risk and egg solls.

References

5-HT2A 5-Hydroxytryptamine Receptor 2A

Camilleri M. Serotonin in the Gastrointestinal Tract. Current opinion in endocrinology, diabetes, and obesity. 2009;16(1):53-59. (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2694720/)

Markoutsaki T, Karantanos T, Gazouli M, Anagnou NP, Karamanolis DG. 5-HT2A receptor gene polymorphisms and irritable bowel syndrome. J Clin Gastroenterol. 2011;45:514–517. (https://www.ncbi.nlm.nih.gov/pubmed/21325954)

ACE Angiotensin I Converting Enzyme

Collins, M.A.L.C.O.L.M., Xenophontos, S.L., Cariolou, M.A., Mokone, G.G., Hudson, D.E., Anastasiades, L. and Noakes, T.D., 2004. The ACE gene and endurance performance during the South African Ironman Triathlons. Medicine and science in sports and exercise, 36(8), pp.1314-1320. (http://europepmc.org/abstract/med/15292738)

Jones N, Kiely J, Suraci B, et al. A genetic-based algorithm for personalized resistance training. Biology of Sport. 2016;33(2):117-126. doi:10.5604/20831862.1198210. (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4885623/)

ACTN3 Alpha 3 Actinin

Eynon, N., Hanson, E.D., Lucia, A. et al. Genes for Elite Power and Sprint Performance: ACTN3 Leads the Way. Sports Med (2013) 43: 803. doi:10.1007/s40279-013-0059-4 (https://www.ncbi.nlm.nih.gov/pubmed/23681449)

Hogarth M.W., Garton F.C, et al; Analysis of the ACTN3 heterozygous genotype suggests that -actinin-3 controls sarcomeric composition and muscle function in a dose-dependent fashion, Human Molecular Genetics, Volume 25, Issue 5, 1 March 2016, Pages 866–877, https://doi.org/10.1093/hmg/ddv613 (https://www.ncbi.nlm.nih.gov/pubmed/26681802)

Yang N, MacArthur DG, Gulbin JP, et al. ACTN3 Genotype Is Associated with Human Elite Athletic Performance. American Journal of Human Genetics. 2003;73(3):627-631. (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1180686/)

ADORA2A Adenosine A2A Receptor

GoldsteinER, ZiegenfussT, KalmanD, et al. (2010) International Society of Sports Nutrition position stand: caffeine and performance. J Int Soc Sports Nutr 7, 5 (https://jissn.biomedcentral.com/articles/10.1186/1550-2783-7-5)

Yang A, Palmer AA, de Wit H. Genetics of caffeine consumption and responses to caffeine. Psychopharmacology. 2010;211(3):245-257. doi:10.1007/s00213-010-1900-1. (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4242593/)

ADRB2 Beta-2-Adrenergic Receptor

Association of the ADRB2 Gly16Arg and Glu27Gln polymorphisms with athlete status Marek Sawczuk, Agnieszka Maciejewska-Karlowska, Pawel Cieszczyk, Bogumila Skotarczak & Krzysztof Ficek Journal of Sports Sciences Vol. 31, Iss. 14,2013 (http://www.tandfonline.com/doi/full/10.1080/02640414.2013.786184?src=recsys)

Lagou V, Liu G, Zhu H, et al. Lifestyle and socioeconomic-status modify the effects of ADRB2 and NOS3 on adiposity in European-American and African-American adolescents. Obesity (Silver Spring) 2011;19:595–603. doi: 10.1038/oby.2010.224. (https://www.ncbi.nlm.nih.gov/pubmed/20930716)

2 Adrenoceptor Functional Gene Variants, Obesity, and Blood Pressure Level Interactions in the General Population Alexandre C. Pereira, Marcilene S. Floriano, Glória F.A. Mota, Roberto S. Cunha, Fernando L. Herkenhoff, José G. Mill, José E. Krueger https://doi.org/10.1161/01.HYP.0000085648.65419.17 Hypertension. 2003;42:685-692 Originally published October 2, 2003 (https://doi.org/10.1161/01.HYP.0000085648.65419.17)

AGT Angiotensinogen

Aleksandra Z, Zbigniew J, Waldemar M, et al. The AGT Gene M235T Polymorphism and Response of Power-Related Variables to Aerobic Training. Journal of Sports Science & Medicine. 2016;15(4):616-624. (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5131215/)

AMPD1 Adenosine Monophosphate Deaminase 1

Ginevičienė V, Jakaitienė A, Pranculis A, Milašius K, Tubelis L, Utkus A. AMPD1 rs17602729 is associated with physical performance of sprint and power in elite Lithuanian athletes. BMC Genetics. 2014;15:58. doi:10.1186/1471-2156-15-58. (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4032451/)

BDKRB2 Bradykinin Receptor Beta 2

Associations of polymorphisms of eight muscle- or metabolism-related genes with performance in Mount Olympus marathon runners Georgios I. Tsianos, Evangelos Evangelou, Arnoud Boot, M. Carola Zillikens, Joyce B. J. van Meurs, Andre G. Uitterlinden, John P. A. Ioannidis Journal of Applied Physiology Mar 2010, 108 (3) 567-574; DOI: 10.1152/japplphysiol.00780.2009 (https://www.ncbi.nlm.nih.gov/pubmed/20044476)

Colleen J. Saunders, Stavroulla L. Xenophontos, Marios A. Cariolou, Lakis C. Anastassiades, Timothy D. Noakes, Malcolm Collins; The bradykinin 2 receptor (BDKRB2) and endothelial nitric oxide synthase 3 (NOS3) genes and endurance performance during Ironman Triathlons. Hum Mol Genet 2006; 15 (6): 979-987. doi: 10.1093/hmg/ddl014 (https://academic.oup.com/hmg/article-lookup/doi/10.1093/hmg/ddl014)

Some pages have been omitted in the sample

Lifecode GX – Professional Genotype Analysis –

Disclaimer

The information provided should not be used for diagnostic or treatment purposes and is not a substitute for personal medical advice. Use the information provided by Lifecode Gx solely at your own risk. Lifecode Gx makes no warranties or representations as to the accuracy of information provided herein. If you have any concerns about your health, please consult a qualified health professional.